

TOTALLY GEODESIC SEIFERT SURFACES IN HYPERBOLIC KNOT AND LINK COMPLEMENTS I

COLIN ADAMS AND ERIC SCHOENFELD

ABSTRACT. The first examples of totally geodesic Seifert surfaces are constructed for hyperbolic knots and links, including both free and totally knotted surfaces. Then it is proved that two bridge knot complements cannot contain totally geodesic orientable surfaces.

1. INTRODUCTION

Define a knot or link in $\mathbf{S}^3$ to be hyperbolic if its complement is a hyperbolic 3-manifold. This implies that there is a covering map p from $\mathbf{H}^3$ to $\mathbf{S}^3 - K$ such that the covering translations are isometries of $\mathbf{H}^3$. We say that an embedded or immersed surface S in $\mathbf{S}^3 - K$ is totally geodesic if it is isotopic to a surface that lifts to a set of geodesic planes in $\mathbf{H}^3$.

Embedded totally geodesic surfaces are relatively rare in knot and link complements. It was conjectured in [MR92] that there are no closed embedded totally geodesic surfaces in hyperbolic knot complements. This has been verified for hyperbolic knots that are alternating knots, tunnel number one knots, and 2-generator knots [MR92], almost alternating knots [Aetal92], toroidally alternating knots [cA94], Montesinos knots [uO84], 3-bridge knots and double torus knots [IO200]) and knots of braid index three [LP85] and four [hM02]. Note that the conjecture does not hold for links. The first explicit counterexample was given in [MR92]. In [Le04], it is demonstrated that for each $g > 2$, there exists a 2-component hyperbolic link that contains a closed embedded totally geodesic genus g surface.

In this paper, we investigate surfaces that are not closed in knot and link complements. In particular, we are interested in totally geodesic Seifert surfaces, where a Seifert surface for a knot or link L is a compact orientable surface whose boundary is L . Note that although a totally geodesic Seifert surface need not necessarily be minimal genus, it is incompressible.

Work of Thurston implies that a properly embedded surface S with boundary in a hyperbolic knot exterior M can have one of three possible behaviors:

- (1) There is a nontrivial simple closed curve c in S that is not boundary parallel in S but that corresponds to a parabolic isometry. This implies that c is isotopic into a neighborhood of the missing knot. In this case, we say that c is an accidental parabolic curve and S is an accidental surface. See [IO200] for more on these surfaces.

- (2) S is a quasi-Fuchsian surface, with limit set a union of quasi-circles. This includes the case of S being totally geodesic (also called Fuchsian), in which case the limit set is a union of geometric circles.
- (3) K is a fibered knot and in some finite cover, S lifts to a fiber. In this case, the limit set is the entire sphere.

In [sF98], it is proved that if K is not fibered, then any minimal genus Seifert surface cannot be accidental and hence must be quasi-Fuchsian. See also [CL01]. A quasi-Fuchsian surface S can admit an essential annuli A intersecting S in its boundary. However, in this case, S cannot be totally geodesic. In essence, the existence of the annulus forces pairs of quasi-circles in the limit set of the cover to share two points, which a pair of circles that do not cross cannot do. So, for a surface to be totally geodesic, it must not admit any essential annuli. Nor can it admit an essential disk with boundary consisting of four arcs, two of which lie in the surface and two of which lie in the cusp. Such a disk doubles to an essential annulus in the manifold obtained by doubling along the totally geodesic surface. However this contradicts the fact the double is a hyperbolic manifold.

A Seifert surface is said to be *free* if the complement of a neighborhood of the Seifert surface is a handlebody. Note that there are examples of knots such that a minimal genus Seifert surface is free and examples such that it is not free. See [yM87].

In Section 2, we explain how lifts of rigid hyperbolic 2-orbifolds can be utilized to generate totally geodesic Seifert surfaces. In Section 3, we utilize these results to generate the first examples of hyperbolic knots with totally geodesic Seifert surfaces. We include both examples of free totally geodesic Seifert surfaces and non-free totally geodesic Seifert surfaces. In Section 4, we prove that many hyperbolic knots cannot possess a totally geodesic Seifert surface, including all two-bridge knots.

Note that the surfaces given here are the only known examples of embedded totally geodesic surfaces of any kind in knot complements in $\mathbf{S}^3$. One can ask whether there are embedded totally geodesic surfaces in knot complements other than Seifert surfaces.

Acknowledgements: Thanks to Satyan Devadoss, Cameron Gordon and Alan Reid for helpful conversations.

2. PRODUCING TOTALLY GEODESIC SEIFERT SURFACES

We call a hyperbolic orbifold *rigid* if there exists only one possible hyperbolic metric that generates the orbifold. In [wT78], it is demonstrated that a hyperbolic 2-orbifold Q with underlying space X_Q , j elliptic points and k corner reflectors is rigid if and only if $3\chi(X_Q) = 2j + k$. A 2-orbifold is hyperbolic if and only if its Euler characteristic is negative where the Euler characteristic is given by:

$$\chi(Q) = \chi(X_Q) - \sum (1 - \frac{1}{m_i}) - \frac{1}{2} \sum (1 - \frac{1}{n_i})$$

where $m_1, m_2, \dots, m_j$ and $n_1, n_2, \dots, n_k$ are the orders of the elliptic points and corner reflectors respectively.

Hence a complete list of rigid hyperbolic 2-orbifolds consists of

- $S^2(m_1, m_2, m_3;)$ for $1/m_1 + 1/m_2 + 1/m_3 < 1$,
- $D^2(; n_1, n_2, n_3)$ for $1/n_1 + 1/n_2 + 1/n_3 < 1$
- $D^2(m_1; n_1)$ for $2/m_1 + 1/n_1 < 1$

Note that each n_i and m_i can be either a positive integer or ∞ .

The fundamental group $\pi_1(O)$ of an orbifold O is the group of deck transformations of the universal cover $\tilde{O}$. A 2-orbifold Q embedded in a 3-orbifold O is *essential* if $\pi_1(Q)$ injects into $\pi_1(O)$. A hyperbolic 2-orbifold Q embedded in a hyperbolic 3-orbifold O is *totally geodesic* if it lifts to the disjoint union of geodesic planes in the universal cover $\mathbf{H}^3$. Let p denote the map projecting $\mathbf{H}^3$ onto O . We state a theorem that follows from the theorem on p. 217 [bM88], but provide a complete proof here for convenience.

Theorem 2.1. *Let Q be an essential rigid hyperbolic 2-orbifold embedded in a hyperbolic 3-orbifold O . Then Q is isotopic relative the singular set to a totally geodesic 2-orbifold Q' .*

Proof. Every isometry γ of hyperbolic 3-space can be represented as the product of two order two elliptic isometries, each a rotation about a geodesic axis. If γ is hyperbolic, then the two axes, considered as subsets of $\mathbf{H}^3 \cup \partial\mathbf{H}^3$, do not intersect. If γ is parabolic then they share an endpoint. If γ is elliptic, then they intersect in $\mathbf{H}^3$ and the axis of rotation for the elliptic isometry passes through the point of intersection and is perpendicular to the plane containing both of the axes.

The fundamental group $\pi_1(O)$ is realized as a discrete group Γ of isometries of $\mathbf{H}^3$ with subgroup Γ' corresponding to $\pi_1(Q)$. There exists a pair of isometries g_1 and g_2 in Γ' , the product of which is a third isometry g_3 in Γ' . In the cases of $S^2(m_1, m_2, m_3;)$ and $D^2(; n_1, n_2, n_3)$, we can choose these isometries corresponding to the three singularities and the three corner reflectors respectively. In the case of $D^2(m_1; n_1)$, we can choose isometries corresponding to the corner reflector and to two lifts of the isometry corresponding to the singularity, one of which is conjugate to the other by an order two elliptic isometry corresponding to the boundary of the orbifold.

Each g_i can be realized as the product of two isometries, each of which is an order two elliptic isometry about an axis. Choose an axis for an order two elliptic isometry f_2 such that if g_1 is elliptic, the axis is perpendicular to the axis of g_1 . If g_1 is parabolic, choose the axis of f_2 so that it ends at the fixed point of g_1 . Also make the axis simultaneously satisfy the same requirements for g_2 . Then we can find order two elliptic isometries f_1 and f_3 such that $g_1 = f_1 f_2$ and $g_2 = f_2 f_3$. Note then that $g_3 = g_1 g_2 = f_1 f_2 f_2 f_3 = f_1 f_3$. Since g_3 is elliptic or parabolic, the axes for f_1 and f_3 must either intersect in $\mathbf{H}^3$ or in $\partial\mathbf{H}^3$. In either case, the three axes for f_1 , f_2 and f_3 all lie in a geodesic plane P' . The entire group Γ' is generated by f_1 , f_2 and f_3 so Γ' preserves P' . The orders of g_1 , g_2 and g_3 determine the angles of the triangle defined by the axes of f_1 , f_2 and f_3 , and hence the resulting 2-dimensional hyperbolic orbifold Q' that is the projection of P' to O is rigid.

Exactly as in the proof of Theorem 3.1 of [cA85], we can prove that the limit set of P' is in fact also the limit set of a topological plane P covering Q . Since Q is embedded, the limit sets of the topological planes covering Q can only intersect tangentially. Hence the same holds for all the copies of P' under the group action, implying that the geodesic planes covering Q' do not intersect in their interiors. In particular, this means that Q' is embedded and $p^{-1}(Q')$ is a disjoint union of geodesic planes, and Q' is a totally geodesic orbifold.

Suppose μ is an element of Γ such that $\mu(P') = P'$. Then μ preserves the limit circle of P' . But then μ preserves the limit set of P , implying $\mu \in \Gamma'$. Hence, Q and Q' have the same fundamental group and are orbifold homeomorphic.

Note that in the case the underlying space of Q' is a disk, Q and Q' must share the boundary arcs, which are order two singularity axes in O .

The orbifold Q can be isotoped so that it does not intersect Q' in its interior as follows. First, put Q in general position relative to Q' . Utilizing the product neighborhoods of the singularities, cusps and boundaries of Q' , we can isotope Q so that there are no intersection curves near singularities, cusps, or boundaries of the two 2-orbifolds. Hence, all the intersection curves are simple closed curves. A trivial intersection curve on one of the 2-orbifolds must be trivial on the other by the fact Q is essential and Q' is totally geodesic. However, such a curve can then be eliminated by isotoping Q .

A curve that is nontrivial on both Q and Q' , and innermost such on Q must bound a disk with one singularity on each of Q and Q' , denoted D and D' respectively. Since this implies that a loop around the singularity on Q is homotopic to the loop around the singularity on Q' , the singularities must either both correspond to parabolic isometries or to elliptic isometries of the same order. Then $D \cup D'$ forms a sphere with two singularities of the same order. By irreducibility, $D \cup D'$ bounds a ball containing an unknotted singular arc or cusp within it. Hence, D can be isotoped through D' to eliminate the intersection.

Now, Q and Q' do not intersect. Hence their lifts, which come paired with shared boundary circles, do not intersect. Thus each pair of planes sharing a boundary circle bound a ball in $\mathbf{H}^3$ that is intersected by none of its translates in the interior. Thus, we can isotope the topological plane that is a lift of Q to the corresponding geodesic plane that is a lift of Q' . In fact, we can do so equivariantly with respect to Γ , and none of the disjoint balls bounded by the planes will intersect in the process. Hence Q is isotopic to Q' , as we wished to demonstrate. $\square$

Corollary 2.2. *Let M be a closed orientable 3-manifold, covering a 3-orbifold O such that M contains a link L covering a collection J of simple closed curves and arcs beginning and ending on the order two singular set of O , and such that $O - J$ is a hyperbolic 3-orbifold O' . Then if O' contains a rigid hyperbolic 2-orbifold Q for which each simple closed component of J is a single puncture (singularity with $m_i = \infty$) and each arc corresponds to a corner reflector with $n_j = \infty$, then Q is isotopic in O' to a 2-orbifold that lifts to a totally geodesic Seifert surface for the link L in M .*

Proof. This follows immediately from Theorem 2.1. $\square$

3. EXAMPLES

In this section, we utilize Corollary 2.2 to give a variety of examples of totally geodesic Seifert surfaces in knot and link complements in $\mathbf{S}^3$.

Example 3.1. *Balanced Pretzel Links*

Let M be $\mathbf{S}^3$ and O the orbifold obtained by quotienting out by a $2\pi/p$ rotation about a great circle C in $\mathbf{S}^3$, where $p \geq 2$. Let D be a disk with two singularities of order p and p , twisted with n crossings as in Figure 1. Let J be the boundary of the disk D . Then J lifts to an $(n, n, n, \dots, n)$ pretzel knot K , where n appears p times. We call K a *balanced pretzel knot*. A picture of the horoball diagram for the $(3, 3, 3)$ pretzel knot appears in Figure 2. This picture was generated by




FIGURE 1. The Seifert surfaces for this family of balanced pretzel links are all totally geodesic.

Jeffrey Weeks' SnapPea computer program, (cf. [jW03]). The sequentially tangent horoballs in a line correspond to a vertical geodesic plane coming up out of the xy -plane that covers the totally geodesic surface.




FIGURE 2. The straight line of horoballs corresponds to the totally geodesic surface.

Example 3.2. *Polyhedral links*

A link that projects to the surface of the cube, as in Figure 3, corresponds to the loop in the orbifold shown, which possesses the shaded rigid 2-orbifold that appears in its complement. Therefore, the link bounds the totally geodesic Seifert surface shown.




FIGURE 3. A totally geodesic Seifert surface for this cube link.

Example 3.3. *Knots with Totally Knotted Seifert Surfaces*

Instead of a disk with two cone points that is twisted relative the singular arc as we saw in Figure 1, we will choose a disk that is knotted relative the singular curve as in Figure 4. When $p = 3$, this lifts to the knot in Figure 4. The disk lifts to a totally geodesic Seifert surface, which is totally knotted, since the complement of a regular neighborhood of the surface is Thurston's wye manifold, which is known to be hyperbolic with incompressible boundary.




FIGURE 4. A totally knotted Seifert surface which is totally geodesic.

4. KNOTS WITHOUT TOTALLY GEODESIC SEIFERT SURFACES

We prove the following theorem.

Theorem 4.1. *Hyperbolic two-bridge knots do not possess orientable totally geodesic surfaces.*

Proof. Note that the only non-hyperbolic 2-bridge knots are the $(2, q)$ -torus knots. There are no non-boundary parallel incompressible closed surfaces in their complements. In [HT85], a complete classification of all the incompressible boundary-incompressible surfaces in the complements of two-bridge knots is given. All such surfaces are carried by branched surfaces that appear as in Figure 5. Such a surface is orientable if and only if each b_i is even where $p/q = r + [b_1, b_2, \dots, b_n]$. For each p/q , there is a unique choice of such even b_i 's.




FIGURE 5. All Seifert surfaces for 2-bridge knots take this basic form.

Note that a Seifert surface corresponds to a 1-sheeted surface carried by the branched surface. Each possible Seifert surface is obtained as a collection of horizontal "plumbing" disks either to the inside or the outside and vertical strips between them, each with an even number of half-twists. The other incompressible boundary-incompressible surfaces have more than one sheet.

If for some $i = 1, 2, \dots, n$, we have $b_i = 2$, then there is a disk in $\mathbf{S}^3$ with boundary that consists of two arcs on S and two arcs on the cusp boundary. When the manifold is doubled along S , this disk doubles to an essential annulus in the resulting manifold. Hence, the resulting manifold cannot be hyperbolic, implying that the surface S could not be totally geodesic.

When $b_i > 2$ for all i , choose a curve c on S as in Figure 5. Note that the curve c can be isotoped to c' as in Figure 5. The curve c' sits on a torus T in $\mathbf{S}^3 - N(S) - N(K)$. Cutting the torus open along c' yields an annulus in $\mathbf{S}^3 - N(S) - N(K)$, both boundary components of which lie on $\partial N(c')$ in T . Gluing on to its boundary components a pair of parallel annuli that give the isotopy from c to c' yields an annulus A in $\mathbf{S}^3 - N(S)$ with boundary in $\partial N(S)$. The annulus A is incompressible in $\mathbf{S}^3 - \text{int}(N(S))$ since its core curve is a power of a generator for the fundamental group of the complementary handlebody. To see that it is boundary-incompressible, we can choose the essential arc α in A shown in Figure 6. If α can be isotoped into the surface, then there exists a disk with boundary consisting of two arcs on S and two arcs on the cusp boundary. Again, when the manifold is doubled along S , this yields an essential annulus, a contradiction to the hyperbolicity of a hyperbolic manifold doubled along a totally geodesic surface within it. If α cannot be isotoped into the surface, then A is an essential annulus intersecting S in its boundary, which obstructs S from being totally geodesic.




FIGURE 6.

□

Do two-bridge knots possess non-orientable totally geodesic surfaces? Any essential non-orientable surface for a 2-bridge knot or link is given by a choice of $[b_1, b_2, \dots, b_n]$, where $p/q = r + [b_1, b_2, \dots, b_n]$ and at least one b_i is odd. The only candidates for totally geodesic surfaces from amongst these are those where each b_i is odd. In the case of links, these surfaces can be totally geodesic. The Whitehead link possesses such a surface. However, we have yet to see such a surface in a 2-bridge knot complement that is totally geodesic.

REFERENCES

- [cA85] Adams, C., “Thrice-punctured spheres in hyperbolic 3-manifolds” *Trans. of AMS* **287** (1985), no. 2, 645–656.
- [Aetal92] Adams, C., Brock, J., Bugbee, J., Comar, T., Faigin, K., Huston, A., Joseph, A., Pesikoff, D., “Almost alternating links” *Topology Appl.* **46** (1992), no. 2, 151–165.
- [cA94] Adams, C., “Toroidally alternating knots and links”, *Topology* **33**, (1994), no. 2, 353–369.
- [CL01] Cooper, D., Long, D.D., “Some surface subgroups survive surgery”, *Geometry and Topology*, **5**, (2001), 347–367.

- [sF98] Fenley, Sergio R., “Quasi-Fuchsian Seifert surfaces” *Math. Z.* 228 (1998), no. 2, 221–227.
- [HT85] Hatcher, A., Thurston, W., “Incompressible surfaces in 2-bridge knot complements” *Invent. Math.* 79 (1985), no. 2, 225–246.
- [IO200] Ichihara, Kazuhiro and Ozawa, Makoto, “Accidental surfaces in knot complements”, *J. Knot Theory Ramifications* 9 (2000), no. 6, 725–733.
- [Le04] Leininger, Christopher, “Small Curvature Surfaces in Hyperbolic 3-Manifolds”, preprint, 2004.
- [LP85] Lozano, M. T., Przytycki, J. H., “Incompressible surfaces in the exterior of a closed 3-braid I”, *Math. Proc. Cambridge Philos. Soc.* 98 (1985), no. 2, 275–299.
- [bM88] Maskit, B., “Kleinian Groups”, Springer-Verlag, Berlin, 1988.
- [hM02] Matsuda, Hiroshi, “Complements of hyperbolic knots of braid index four contain no closed embedded totally geodesic surfaces”, *Topology Appl.* 119(2002), no. 1, 1–15.
- [MR92] Menasco, William and Reid, Alan, “Totally geodesic surfaces in hyperbolic link complements”, *Topology '90 (Columbus, OH, 1990)*, 215–226, Ohio State Univ. Math. Res. Inst. Publ., 1, de Gruyter, Berlin, 1992.
- [yM87] Moriah, Yoav, “On the free genus of knots”, *Proc. Amer. Math. Soc.* 99 (1987), no. 2, 373–379.
- [wT78] Thurston, William, *The geometry and topology of 3-manifolds*, Lecture Notes, 1978.
- [uO84] Oertel, Ulrich, “Closed incompressible surfaces in complements of star links”, *Pac. J. of Math.*, Vol. 111, No. 1, 1984, 209–230.
- [jW03] Weeks, Jeffrey, “Snappea Computer program”, available at www.geometrygames.org.

COLIN ADAMS, DEPARTMENT OF MATHEMATICS AND STATISTICS, WILLIAMS COLLEGE, WILLIAMSTOWN, MA 01267

E-mail address: `Colin.C.Adams@williams.edu`

ERIC SCHOENFELD, DEPARTMENT OF MATHEMATICS, STANFORD UNIVERSITY, 450 SERRA MALL, BLDG. 380, STANFORD, CA 94305-2125

E-mail address: `erics@math.stanford.edu`